JPL-IN-418-CR 326640 P.58

JPL Publication 90-40

Bio-Optical Profile Data Report

Coastal Transition Zone Program R/V Thomas Washington June 24–July 21, 1988

Curtiss O. Davis W. Joseph Rhea

December 1, 1990

Prepared for

Office of Naval Research
Through an agreement with

National Aeronautics and Space Administration

by

Jet Propulsion Laboratory California Institute of Technology Pasadena, California

(NASA-CR-187808) DIO-OPTICAL PROFILE DATA REPORT COASTAL TRANSITION ZONE PROGRAM, R/V THOMAS WASHINGTON, JUNE 24 - JULY 21, 1988 (JPL): 58 p CSCL 08C

N91-16532

Unclas G3/48 0326640 4. 🖍

•	TEC	HNICAL REPORT STANDARD TITLE PAGE	
1. Report No. JPL Pub. 90-40	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle	5. Report Date December 1, 1990		
Bio-Optical Profile Data Re Coastal Transition Zone Pro R/V Thomas Washington Ju	6. Performing Organization Code		
7. Author(s) Custiss O. Davis a	8. Performing Organization Report No. JPL Publication 90-40		
9. Performing Organization Name ar	10. Work Unit No.		
JET PROPULSION LAB California Institu 4800 Oak Grove Dri	11. Contract or Grant No. NAS7-918		
Pasadena, California 91109		13. Type of Report and Period Covered	
12. Sponsoring Agency Name and Ad	dress	JPL Publication	
NATIONAL AERONAUTICS AND Washington, D.C. 20546	14. Sponsoring Agency Code		
15. Supplementary Notes Prepared for Office of Nav Aeronautics and Space Admin	ral Research through an agraistration by the Jet Propu	reement with the National	
16. Abstract			
1			

Twenty-three vertical profiles of the bio-optical properties of the ocean were made during a research cruise on the R/V Thomas Washington, June 24--July 21, 1988, as part of the Coastal Transition Zone Program off Point Arena, California. This report is a summary to provide investigators with an overview of the data collected. The entire data set is available in digital form for interested researchers, and requests for the data should be addressed to W. Joe Rhea, (818) 393-6095.

17. Key Words (Selected by Author(s)	18. Distribution Statement			
Geosciences and Oceanogra Biological Oceanography	phy (general)	Unlimited/Unc	lassified	
19. Security Classif. (of this report)	20. Security C	lassif. (of this page)	21. No. of Pages	22. Price
Unclassified	Unclass	ified	56	

÷			- 100

Bio-Optical Profile Data Report

Coastal Transition Zone Program R/V Thomas Washington June 24–July 21, 1988

Curtiss O. Davis W. Joseph Rhea

December 1, 1990

Prepared for

Office of Naval Research
Through an agreement with

National Aeronautics and Space Administration

by

Jet Propulsion Laboratory California Institute of Technology Pasadena, California The research described in this publication was carried out by the Jet Propulsion Laboratory, California Institute of Technology, and was sponsored by the Office of Naval Research through an agreement with the National Aeronautics and Space Administration.

Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not constitute or imply its endorsement by the United States Government or the Jet Propulsion Laboratory, California Institute of Technology.

ABSTRACT

Twenty-three vertical profiles of the bio-optical properties of the ocean were made during a research cruise on the R/V Thomas Washington, June 24–July 21, 1988, as part of the Coastal Transition Zone Program off Point Arena, California. This report is a summary to provide investigators with an overview of the data collected. The entire data set is available in digital form for interested researchers, and requests for the data should be addressed to W. Joe Rhea, (818) 393-6095.

ACKNOWLEDGMENTS

The assistance of the Captain and crew of the R/V Thomas Washington and Chief Scientist Tim Cowles is gratefully acknowledged.

CONTENTS

INTRO	DUC'	TION .			1
DATA	DES	CRIPTI	ON		1
INDIV	IDUA	L STAT	ION DATA	PROFILES	1
REFE	RENC	ES			3
TABLE	ES				
	I. II.	Data C Station	Channels Summary		4 5
FIGUR	ES.				
	1.	File:	c880704c		6
	2.	File:	c880705c	***************************************	
	3.	File:	c880705d	***************************************	
	4.	File:	c880706b		12
	5.	File:	c880706c		14
	5. 6.	File:	c880707a		16
		File:	c880707a		
	7.		c8807070		20
	8.	File:			20
	9.	File:	c880708b		21
	10.	File:	c880709b	•••••	24
	11.	File:	c880710a		20
	12.	File:	c880710b		20 20
	13.	File:	c880711b		20
	14.	File:	c880714a		34
	15.	File:	c880715a		
	16.	File:	c880716b		36
	17.	File:	c880717a		38
	18.	File:	c880718a		40
	19.	File:	c880719a	***************************************	42
	20.	File:	c880720a	***************************************	44
	21.	File:	c880720c	***************************************	46
	22.	File:	c880720e	***************************************	48
	23	File:	c880721a		50

INTRODUCTION

The Coastal Transition Zone (CTZ) Program, sponsored by the Office of Naval Research, is an interdisciplinary study of the physical causes and the physical, biological, and optical characteristics of the filaments of cold, salty water that extend over 300 km off the West Coast of North America. The cruise of the R/V Thomas Washington, June 24–July 21, 1988, was designed to study a representative filament off Point Arena, California. Guided by satellite sea-surface temperature maps, two surveys of the filament and adjacent waters were conducted. Additionally, two clusters of drifters were placed in the root of the filament and tracked by satellite for several months. One of the drifters in the first cluster was instrumented with a fluorometer, transmissometer, spectroradiometer, and a water sampler. At the end of seven days, the instrumented drifter was recovered approximately 300 km to the southwest of the launch site.

This report describes bio-optical profile data that were collected at 23 stations representative of the filament, freshly upwelled water near the coast, and the offshore water surrounding the filament. Of particular interest is a time series of seven stations taken at approximately noon each day next to the instrumented drifter.

DATA DESCRIPTION

Optical data were collected with a Bio-Optical Profiling System (BOPS), an updated version of the BOPS originally developed by Smith et al. (1984). The heart of the BOPS is a Biospherical Instruments MER-1048 spectroradiometer, which measured up- and downwelling spectral irradiance and upwelling spectral radiance. The MER-1048 also has sensors for Photosynthetically Available Radiation (PAR), depth, tilt, and roll. In addition, temperature and conductivity were measured with a Sea-Bird CTD, chlorophyll fluorescence was measured with a Sea Tech fluorometer, and beam transmission was measured with a Sea Tech 25-cm transmissometer. The MER-1048 acquired all the data 16 times a second, averaged it to four records a second, and sent it up the cable to a deck box and a Compaq-286 computer, which stored the data on the hard disk. The BOPS data (Table I) were filtered to remove obvious data spikes and then binned into one-meter averages and stored in the form of ASCII comma-separated files.

INDIVIDUAL STATION DATA PROFILES

For each station (Table II), eight profiles are presented to give a graphical overview of the data (Figures 1–23). Data files are identified by a filename of the format:

Nyymmddc.MER

```
where:
N = c for CTZ cruise of the R/V Thomas Washington, June 24–July 21, 1987
yy = Year
mm = Month
dd = Day
c = Cast order for each day.
i.e. 'a' = first cast of day
'b' = second cast of day, etc.
```

Temperature and salinity data were from the Sea-Bird CTD. Salinity was calculated from the temperature and conductivity measurements using the standard equations for practical salinity units (Millero et al., 1980). Occasional spikes in salinity were observed at the

surface and at the thermocline. This is an artifact caused by the fact that the response time of the conductivity sensor does not exactly match that of the temperature sensor.

Data from the Sea Tech fluorometer are presented in fluorescence units. The fluorometer data were calibrated using extracted chlorophyll and phaeopigment values from water samples taken immediately before or after a number of optical profiles. Average chlorophyll plus phaeopigment values for the entire cruise give the following equation for calibrating the Sea Tech fluorometer data: chl + phaeo = 0.288 + 0.218 fluor, where $r^2 = 0.77$ and n = 136.

Beam transmissometer (25-cm path length, 660-nm wavelength) data were recorded in percent transmission (%T; value in air was set to 100%). The attenuation of a beam of light is defined by Jerlov (1976):

c = a + b

and

 $T = e^{-cr}$

where

c is the beam attenuation coefficient in m⁻¹

a is the absorption coefficient

b is the total scattering coefficient

T is the fraction of light transmitted over path length r.

Then, for this data set, c can be calculated from the following equation (r = 0.25 m):

$$c = -4 ((\%T/100)*.949)$$

The radiance, irradiance, and PAR data are presented in calibrated units based on a laboratory calibration conducted by Biospherical Instruments on June 21, 1988. A second calibration after the cruise showed no significant deviation from these values. The spectral light data are presented as a plot of spectra near the surface (dashed line) and then at every five meters (5 m, 10 m, 15 m, etc.). Typically, the surface reading is for 2 m; however, during rough weather, the first usable readings are from greater depth, as indicated on the figures. This is calibrated radiance data, but no corrections for ship shadow or other artifacts have been made to the data. We have developed routines for correcting such artifacts, calculating K, etc., following the guidelines of Smith and Baker (1984, 1986) and Gordon (1985), and the reader is referred to those references for a discussion of these problems.

The data are available in digital format for researchers who wish to work with the actual data. Individuals who are interested in working with the data should request it in digital form from Joe Rhea ((818) 393-6095). The data can be provided in a number of formats compatible with most standard computing environments.

REFERENCES

- Gordon, H. R., "Ship Perturbation of Irradiance Measurements at Sea. 1: Monte Carlo Simulations," *Applied Optics*, 24, 4172-4182, 1985.
- Jerlov, N. G., Marine Optics, Elsevier, New York, 1976.
- Millero, F., C.-T. Chien, A. Bradshaw, and K. Schleicher, "A New High Pressure Equation of State for Seawater," *Deep-Sea Research*, 27a, 255-264, 1980.
- Smith, R. C., and K. S. Baker, "The Analysis of Ocean Optical Data," Proceedings of the SPIE, Ocean Optics VII, 489, 119-126, 1984.
- Smith, R. C., and K. S. Baker, "Analysis of Ocean Optical Data II," Proceedings of the SPIE, Ocean Optics VIII, 637, 95-107, 1986.
- Smith, R. C., C. R. Booth, and J. L. Star, "Oceanographic Bio-optical Profiling System," *Applied Optics*, 23, 2791-2797, 1984.

TABLE I.


Data Channels


- 0. Number of data points averaged into bin
- 1. 410-nm Downwelling Irradiance (µW/cm²/nm)
- 2. 441-nm Downwelling Irradiance (µW/cm²/nm)
- 3. 488-nm Downwelling Irradiance (μW/cm²/nm)
- 4. 520-nm Downwelling Irradiance (μW/cm²/nm)
- 5. 550-nm Downwelling Irradiance (μW/cm²/nm)
- 6. 560-nm Downwelling Irradiance (μW/cm²/nm)
- 7. 589-nm Downwelling Irradiance (µW/cm²/nm)
- 8. 633-nm Downwelling Irradiance (µW/cm²/nm)
- 9. 656-nm Downwelling Irradiance (µW/cm²/nm)
- 10. 671-nm Downwelling Irradiance (μW/cm²/nm)
- 11. 683-nm Downwelling Irradiance (μW/cm²/nm)
- 12. 694-nm Downwelling Irradiance (μW/cm²/nm)
- 13. 710-nm Downwelling Irradiance (µW/cm²/nm)
- 14. Depth of averaged bin (m)
- 15. Tilt (angles in degrees (-45 to +45))
- 16. Roll (angles in degrees (-45 to +45))
- 17. 410-nm Radiance (μW/cm²/nm/sr)
- 18. 441-nm Radiance (μW/cm²/nm/sr)
- 19. 488-nm Radiance (μ W/cm²/nm/sr)
- 20. 520-nm Radiance (μW/cm²/nm/sr)
- 21. 550-nm Radiance (μ W/cm²/nm/sr)
- 22. 633-nm Radiance (µW/cm²/nm/sr)
- 23. 656-nm Radiance (μW/cm²/nm/sr)
- 24. 683-nm Radiance (µW/cm²/nm/sr)
- 25. 410-nm Upwelling Irradiance (μW/cm²/nm)
- 26. 441-nm Upwelling Irradiance (μW/cm²/nm)
- 27. 488-nm Upwelling Irradiance (μW/cm²/nm)
- 28. 520-nm Upwelling Irradiance (μW/cm²/nm)
- 29. 550-nm Upwelling Irradiance (μW/cm²/nm)
- 30. 589-nm Upwelling Irradiance (μW/cm²/nm)
- 31. 671-nm Upwelling Irradiance (μ W/cm²/nm)
- 32. 694-nm Upwelling Irradiance (µW/cm²/nm)
- 33. Transmissometer 25 cm (% transmission)
- 34. Fluorometer (fluorescence units)
- 35. PAR (10¹⁷quanta/cm²/s]
- 36. Temperature (deg C)
- 37. Conductivity (mmho/cm)
- 38. Salinity (PSU)
- 39. Density (g/cm³)
- 41. 520-nm Surface Irradiance (ship mounted)
- 40. 410-nm Surface Irradiance (ship mounted)
- 42. 589-nm Surface Irradiance (ship mounted)
- 43. 683-nm Surface Irradiance (ship mounted)


TABLE II.


1988 R/V Thomas Washington CTZ Cruise–Station Summary


FILENAME	Sta	Time	Latitude	Longitude	Comments
yymmdd	(Cast)	(Ship)	Deg Min	Deg Min	
c880704c c880705c c880705d c880706b c880706c c880707a c880707b c880708b c880709b c880710a c880710b c880711b c880715a c880715a c880717a c880717a c880717a c880719a c880720a c880720c c880720e c880721a	1(1) 5(2) 6(1) 10(2) 11(2) 16(2) 17(2) 21(2) 22(2) 27(2) 31(2) 32(1) 37(3) 52(2) 63(2) 66(2) 70(1) 72(2) 77(2) 80(2) 82(1) 84(1) 91(1)	15:41 08:01 10:40 07:57 11:32 07:29 14:45 07:22 12:41 12:46 07:28 11:30 10:07 11:02 11:25 12:09 11:05 10:57 09:01 12:29 16:51 16:09	N39 15.0 N38 55.1 N38 49.8 N38 20.3 N38 24.2 N38 06.6 N38 07.2 N37 45.5 N37 53.5 N37 29.7 N37 07.6 N37 16.2 N37 33.3 N38 43.4 N38 37.8 N37 44.7 N37 36.2 N37 14.2 N38 19.7 N37 37.1 N37 45.8 N37 64.3 N38 57.6	W124 13.1 W124 10.9 W124 20.1 W124 09.2 W124 22.3 W124 34.8 W124 54.1 W125 03.6 W125 37.9 W126 02.3 W126 09.7 W126 32.1 W126 34.4 W124 00.0 W124 33.2 W125 29.8 W126 23.7 W127 15.5 W126 44.5 W124 28.6 W124 36.9 W124 57.7	Drifter #1 South of drifter Drifter #2 South of drifter Drifter #3, C-14 prod South of drifter Drifter #4, C-14 prod South of drifter Drifter #5, C-14 prod Drifter #6, C-14 prod South of drifter Drifter #7 Blue water In nearshore eddy West of jet, C-14 prod 2nd drifter, C-14 prod 2nd drifter, C-14 prod 2nd drifter, C-14 prod South of jet, D-line In jet, D-line, C-14 prod North of jet, D-line






































































































































		•

		_
		•
	•	
